

85. (Twice amended) An assay for determining the presence or amount of a free and complexed cardiac specific isoform of troponin in a patient sample, said assay comprising:

obtaining a sample; adding to the sample
[performing an immunoassay with] an antibody which specifically binds said free cardiac specific isoform of troponin, and which specifically binds said cardiac specific isoform of troponin in a binary complex comprising one other troponin component selected from the group consisting of troponin I, troponin C and troponin T, and which specifically binds said cardiac specific isoform of troponin in a ternary complex comprising two other troponin components selected from the group consisting of troponin I, troponin C and troponin T; and

C1
detecting a signal [from said immunoassay] resulting from said antibody binding said free and complexed cardiac specific isoform of troponin, wherein said signal is at least a factor of two larger than a signal resulting from said antibody binding to an equal number of (i) free troponin components which are not said cardiac specific isoform of troponin; (ii) troponin complexes which do not comprise said cardiac specific isoform of troponin; or (iii) a combination of (i) and (ii), and wherein said signal is related to the presence or amount of said free and complexed cardiac specific isoform of troponin in said sample.

88. (Twice amended) An assay for determining the presence or amount of a free and complexed cardiac specific isoform of troponin in a patient sample, said assay comprising:

C2
performing an immunoassay with an antibody which specifically binds to free cardiac specific troponin I and cardiac specific troponin T, and which specifically binds to cardiac specific troponin I and cardiac specific troponin T in a complex comprising at least one other troponin component selected from the group consisting of troponin I, troponin C and troponin T, and which specifically binds to cardiac specific troponin I and cardiac specific troponin T in a ternary complex comprising two other troponin components selected from the group consisting of troponin I, troponin C and troponin T; and

C2
cond
detecting a signal from said immunoassay resulting from said antibody binding said free and complexed cardiac specific isoform of troponin, wherein said signal is at least a factor of two larger than a minimum signal resulting from said antibody binding to an equal number of (i) free troponin components which are not said cardiac specific isoform of troponin; (ii) troponin complexes which do not comprise said cardiac specific isoform of troponin; or (iii) a combination of (i) and (ii), and wherein said signal is related to the presence or amount of said free and complexed cardiac specific isoform of troponin in said sample.

91. (Twice amended) An assay for determining the presence or amount of free and complexed cardiac specific troponin I in a patient sample, said assay comprising:

performing an immunoassay with an antibody which specifically binds to free cardiac specific troponin I, and which specifically binds to cardiac specific troponin I in a ^{binary} complex comprising at least one other troponin component selected from the group consisting of troponin C and troponin T, and which specifically binds to cardiac specific troponin I in a ternary complex comprising troponin C and troponin T; and

C3
detecting a signal from said immunoassay resulting from said antibody binding said free and complexed cardiac specific troponin I, wherein said signal is at least a factor of two larger than a signal resulting from said antibody binding to an equal number of (i) free troponin components which are not said cardiac specific troponin I; (ii) troponin complexes which do not comprise said cardiac specific troponin I; or (iii) a combination of (i) and (ii), and wherein said detectable signal is related to the presence or amount of said free and complexed cardiac specific troponin I in said sample.

94. (Twice amended) An assay for determining the presence or amount of free and complexed cardiac specific troponin T in a patient sample, said assay comprising:

CA
performing an immunoassay with an antibody which specifically binds to free cardiac specific troponin T, and which specifically binds to cardiac specific troponin T in a complex

comprising at least one other troponin component selected from the group consisting of troponin C and troponin I, and which specifically binds to cardiac specific troponin T in a ternary complex comprising troponin C and troponin I; and

C4
COND
detecting a signal from said immunoassay resulting from said antibody binding said free and complexed cardiac specific troponin T, wherein said signal is at least a factor of two larger than a signal resulting from said antibody binding to an equal number of (i) free troponin components which are not said cardiac specific troponin T; (ii) troponin complexes which do not comprise said cardiac specific troponin T; or (iii) a combination of (i) and (ii), and wherein said detectable signal is related to the presence or amount of said free and complexed cardiac specific troponin T in said sample.

102. (Twice amended) An assay for determining the presence or amount of all free and complexed cardiac specific isoform of troponin in a patient sample, said assay comprising:

C5
performing an immunoassay with an antibody which specifically binds all free cardiac specific isoforms of troponin, and which specifically binds all cardiac specific isoforms of troponin in a complex comprising at least one other troponin component selected from the group consisting of troponin I, troponin C and troponin T, and which specifically binds all cardiac specific isoforms of troponin in a ternary complex comprising two other troponin components selected from the group consisting of troponin I, troponin C and troponin T; and

detecting a signal from said immunoassay resulting from said antibody binding said free and complexed cardiac specific isoforms of troponin, wherein said signal is related to the presence or amount of all free and complexed cardiac specific isoform of troponin in said sample.

C6
114. An assay for determining the presence or amount of a free and complexed cardiac specific isoform of troponin in a patient sample, said assay comprising:

Cle
Cans
performing an immunoassay with an antibody which specifically binds said free cardiac specific isoform of troponin, and which specifically binds said cardiac specific isoform of troponin in a complex comprising at least one other troponin component selected from the group consisting of troponin I, troponin C and troponin T, and which specifically binds said cardiac specific isoform of troponin in a ternary complex comprising two other troponin components selected from the group consisting of troponin I, troponin C and troponin T; and

detecting a signal from said immunoassay resulting from said antibody binding said free and complexed cardiac specific isoform of troponin, wherein said signal is related to the presence or amount of said free and complexed cardiac specific isoform of troponin in said sample.

119. (Amended) An assay for determining the presence or amount of a free and complexed cardiac specific isoform of troponin in a patient sample, said assay comprising:

Cut
DL
C1
performing an immunoassay with an antibody which specifically binds to free cardiac specific troponin I and cardiac specific troponin T, and which specifically binds to cardiac specific troponin I and cardiac specific troponin T in a complex comprising at least one other troponin component selected from the group consisting of troponin I, troponin C and troponin T, and which specifically binds to cardiac specific troponin I and cardiac specific troponin T in a ternary complex comprising two other troponin components selected from the group consisting of troponin I, troponin C and troponin T; and

detecting a signal from said immunoassay resulting from said antibody binding said free and complexed cardiac specific isoform of troponin, wherein said signal is related to the presence or amount of said free and complexed cardiac specific isoform of troponin in said sample.